

Electric Blanket Use and Breast Cancer

To the Editor:

Those involved in the Electromagnetic Fields and Breast Cancer on Long Island Study (EBCLIS)^{1,2} have reported significant and conclusive findings in the long-running debate of an EMF–breast cancer relationship. Because the EBCLIS used residential EMF measures and considered characteristics related to proposed mechanisms of causation (eg, stratifying by menopause status and use of hormone receptor status), this study has a degree of precision compared with similar studies, which have produced largely inconsistent results. Kabat and colleagues¹ have investigated a broad range of measures of electromagnetic field exposures and electric blanket use, and the thoroughness of the study unexpectedly gives rise to a number of questions about the informativeness of some of the variables studied. EBCLIS has provided evidence that risk of breast cancer resulting from EMF will not be revealed using the variables and measures reported. A fundamentally important EBCLIS outcome and an apparent measure of EMF from electric blankets have been left out of consideration.

EBCLIS authors reported no differences between cases and controls in all residential EMF exposure measurements. Although these variables were appropriately not included in the authors' regression model, given the lack of difference in residential EMF fields between cases and controls, the finding of no association between electric blanket use and breast cancer risk could not be surprising.¹ Any effect of EMF fields from electric blankets would need to be effectively larger than that of residential EMF levels; otherwise, background EMF levels might mask the effects of electric blankets. Another possibility is that the effect of electric blanket EMF was buried in the proxy, self-reported measures used to estimate the magnitude of EMF effect and could well have

been uncovered if direct measures of EMF strength from electric blankets were taken. Any similarities or differences between cases and controls would be realized immediately by taking quantitative measures of EMF directly from study subjects' blankets. Because EMF is central to the EBCLIS author's hypothesis, these measures could have provided knowledge of the possible success of finding an association. To this end, the authors overlooked an important measure of electric blanket use despite their comprehensive number of variables.

In the study of electric blankets,¹ the authors included a novel variable to test, ie, whether the study subject's windows were opened or closed during the night. They state, and truly so, that this should increase electric blanket cycling through the night, because the blanket tries to maintain a set temperature. However, has this possible effect been accurately modeled? Although the authors presented this intriguing possibility, there is no mention of its reality. If evidence of increased risk of breast cancer with this variable were found, would the study then revisit the phenomenon to see if it exists? This effect could have been modeled appropriately using electric blanket measures and could have provided further strength to their model. Thus, direct measures of electric blanket EMF could have also provided insight into the window open/closed variable considered.

This study¹ provides evidence of no risk of breast cancer resulting from EMF or electric blanket use, or at least convincing support of the fact that any association between the two will not be realized with the methods used. Although the study is comprehensive, the authors need to address a number of basic questions regarding possible differences in residential EMF levels between cases and controls, and they should consider the use of direct measures of blanket EMF levels. There is need for more direct and increasingly quantitative measures to tease out any

possibly concealed relationship between EMF and breast cancer.

Jason Robarge

Department of Human Genetics
Washington University in St. Louis
St. Louis, MO
jrobarge@wubios.wustl.edu

REFERENCES

1. Kabat GC, O'Leary ES, Schoenfeld ER, et al. for the EBCLIS Group. Electric Blanket Use and Breast Cancer on Long Island. *Epidemiology*. 2003;14:514–520.
2. Schoenfeld ER, O'Leary ES, Henderson K, et al. for the EBCLIS group. Electromagnetic fields and breast cancer on Long Island: a case-control study. *Am J Epidemiol*. 2003;158:47–58.

The authors respond:

We thank Jason Robarge¹ for his letter regarding our article.² Although direct measurements of the electric blankets actually used by the participants might be of interest, such measurements do not necessarily provide a valid assessment of EMF exposure and were not feasible in our study for the reasons that follow.

1. Because our study had a case-control study design and ascertained the past use of electric blankets, it is likely that participants used several blankets over their lifetime. Many of the older blankets would have been discarded and not available for measurements.
2. Furthermore, most of the participants who used electric blankets were former users (77%); as such, their blankets would not have been in use during the study period.
3. Even for current users, blanket measurements would not provide an indication of past exposure. Before 1989, there were 3 blanket manufacturers, and exposure levels produced by these earlier or conventional blankets were higher than in later years.³ Newer blankets were manufactured using the PTC (positive temperature coefficient) design, which allowed for cancellation of fields and much lower levels of magnetic fields emit-

ted.⁴ Most participants who used electric blankets purchased their blankets before 1989 (88%). We found no association when stratifying by use of older blankets or newer blankets, although the number of participants who reported using the newer blankets was small.

4. We interviewed participants during all 4 seasons of the year. Valid measurements of the electric blankets as they were actually used (eg, use when the window was open or closed) would be difficult. Our results are consistent with the findings of 8 previous studies of electric blankets and breast cancer.

Erin S. O'Leary

Department of Preventive Medicine
School of Medicine
Stony Brook University
Stony Brook, NY
eoleary@notes.cc.sunysb.edu

Geoffrey C. Kabat*
Elinor R. Schoenfeld
M. Cristina Leske

Department of Preventive Medicine
Stony Brook University
Stony Brook, NY

*Affiliation at the time the research was done.

REFERENCES

1. Robarge J. Electric blanket use and breast cancer [Letter]. *Epidemiology*. 2004;15:375.
2. Kabat GC, O'Leary ES, Schoenfeld ER, et al. for the EBCLIS group. Electric blanket use and breast cancer on Long Island. *Epidemiology*. 2003;14:514–520.
3. Florig HK, Hoburg JF. Power frequency magnetic fields from electric blankets. *Health Physics*. 1990;58:493–502.
4. US Environmental Protection Agency. *EMF in Your Environment: Magnetic Field Measurements of Everyday Electrical Devices*. Washington, DC: Office of Radiation and Indoor Air; 1992 (Publication no. 405-R-92-008).

To the Editor:

A recent editorial¹ asserts that the study by Kabat and colleagues² has done “as well as an epidemiologic design is capable of doing” in assessing the effect of electric blanket use on breast cancer.

However, I believe that the study has a number of severe limitations that render it unlikely the “intriguing hypothesis” of a contribution of exposure to electromagnetic fields (EMFs) by electric blanket use to the risk of breast cancer “can now be put to rest,” as the editorial concludes. A study considered to be “negative” has to show at least that 1) the design is compatible with the study hypothesis, 2) the observed effect was below a margin derived from considerations of the mechanism of action, and 3) the power was sufficient to detect an effect at that level. The study lacks all 3 prerequisites.

The authors start from the hypothesis that extremely low-frequency (ELF) EMF could reduce nocturnal melatonin levels, which could in turn lead to elevated estrogen concentrations, thereby increasing breast cancer risk.³ Although bright light at night suppresses melatonin, there is only equivocal evidence for ELF–EMF to produce such an effect. It has been indicated recently⁴ that exposure to ELF magnetic fields increases cell proliferation in the mammary epithelium without affecting pineal or tissue levels of melatonin. Some insight into the action of ELF–EMFs could be gained from the observation^{5,6} that ELF–EMFs reduce the inhibitory effect of melatonin on cell growth in breast cancer cells. Hence, the melatonin hypothesis could be reformulated as follows. Exposure inhibits antimitogenic pathways, in particular the action of melatonin, leading to increased turnover rates in either stem cells or precancerous lesions thereby increasing the probability of malignant transformation or reducing latency of manifest disease.

If a promotional effect is assumed, then the design and analysis of this study² are inadequate. Because a promoter that acts for a relatively short duration of time only shifts the distribution of age at diagnosis, it either does not affect the overall incidence or only slightly increases it. Clearly, reduction of latency cannot exceed a fraction of exposure duration. Median duration of

use in the EBCLIS study was approximately 6 years. Because controls were frequency-matched by age in 5-year strata, any effect must have been greatly attenuated. Even if age at onset were shortened by 50% of exposure duration, incidence by age stratum would remain almost the same. Effects of this type could be studied by cohort approaches applying multistage models of carcinogenicity, as has been done previously for breast cancer.⁷

A further weakness of the article² is its lack of any consideration regarding the magnitude of the effect of electric blanket use. If the study hypothesis is true, then we seek an effect size that is compatible with what is known about effects of circulating estrogens and progestins on breast cancer risk. A reanalysis of 51 epidemiologic studies of hormone replacement therapy (HRT) on breast cancer risk⁸ reported an average relative risk of 1.35. Recently published hazard ratios from large randomized, controlled trials of HRT^{9,10} were 1.26 and 1.27, with a noticeable increase starting approximately 4 years after onset of therapy. Thus, the hypothesized indirect effect of EMFs is not compatible with effects higher than those mirrored in odds ratios of 1.1 to 1.2. Furthermore, an exposure duration of at least 4 years has to be allowed for. Because at present there is evidence for a reversal of the effect of elevated hormone levels, cases with cessation of exposure approximately 5 years before reference date must be considered unexposed. Given these conditions, both the EBCLIS and the Long Island Breast Cancer Study Project have a negligibly small power to detect an effect of electric blanket use.

Michael Kundi

Institute of Environmental Health
University of Vienna
Vienna, Austria
michael.kundi@univie.ac.at

REFERENCES

1. Sandler DP. On electric blankets and breast cancer [Editorial]. *Epidemiology*. 2003;14:509.
2. Kabat GC, O'Leary ES, Schoenfeld ER, et al. Electric blanket use and breast cancer on Long Island. *Epidemiology*. 2003;14:514–520.
3. Stevens RG. Electric power use and breast cancer: a hypothesis. *Am J Epidemiol*. 1987;125:556–561.
4. Fedrowitz M, Westermann J, Loeschner W. Magnetic field exposure increases cell proliferation but does not affect melatonin levels in the mammary gland of female Sprague Dawley rats. *Cancer Res*. 2002;62:1356–1363.
5. Liburdy RP, Sloma TS, Sokolic R, et al. ELF magnetic fields, breast cancer, and melatonin: 60 Hz fields block melatonin's oncostatic action on ER+ breast cancer cell proliferation. *J Pineal Res*. 1993;14:89–97.
6. Blackman CF, Benane SG, House DE. The influence of 1.2 μ T, 60 Hz magnetic fields on melatonin- and tamoxifen-induced inhibition of MCF-7 cell growth. *Bioelectromagnetics*. 2001;22:122–128.
7. Colditz GA, Rosner B. Cumulative risk of breast cancer to age 70 years according to risk factor status: data from the Nurses' Health Study. *Am J Epidemiol*. 2000;152:950–964.
8. Beral V for the Collaborative Group on Hormonal Factors in Breast Cancer. Breast cancer and hormone replacement therapy: collaborative reanalysis of data from 51 epidemiological studies of 52,705 women with breast cancer and 108,411 women without breast cancer. *Lancet*. 1997;350:1047–1059.
9. Writing Group for the Women's Health Initiative Investigators. Risks and benefits of estrogen plus progestin in healthy postmenopausal women: principal results from the Women's Health Initiative randomized controlled trial. *JAMA*. 2002;288:321–333.
10. Hulley S, Furberg C, Barrett-Connor E, et al. for the HERS Research Group. Noncardiovascular disease outcomes during 6.8 years of hormone therapy. Heart and Estrogen/Progestin Replacement Study follow-up (HERS II). *JAMA*. 2002;288:58–66.

The authors respond:

Dr. Kundi¹ raises concerns about the ability of our study² to detect an association between the use of electric blankets and breast cancer, on several grounds. The first is that ours was not a cohort study; the second relates to the small magnitude of the expected effect; and the third is that, despite our population-based case-control sample size of 2780 women, power was limited to formally evaluate whether recent electric blanket use or long duration of use increases a woman's risk of breast cancer.

In fact, our study was specifically designed to test the hypothesis that ever-use of electric blankets was associated with increased risk of breast cancer, and we had sufficient power to address this question. While cohort studies may be ideal, case-control studies can also provide valuable insight. As to the small magnitude of effect that Kundi projects, individual epidemiologic studies are rarely capable of reliably detecting relative risks on the order of 1.1 to 1.2. We address Kundi's points in light of our results and those of other studies.

We collected detailed information on lifetime use of electric blankets to explore whether various aspects of exposure influence breast cancer development. In contrast to finding an increased risk, the odds ratios for subgroups of ever-users, adjusted for potential confounders, suggest a reduction in both pre- and postmenopausal breast cancer risk, both in current users and in users for more than 10 years. However, the confidence intervals for these estimates cannot rule out a very modest increase in risk. Thus, as with all observational studies, our results need to be interpreted in the context of previous studies on this question. None of the 8 studies we cited previously,² including the Nurses' Health cohort study,³ observed a statistically significant association between electric blanket use and breast cancer. Furthermore, it is striking that none of the studies that looked at duration of use showed any hint of an association, as one would expect for a promoter. Since publication of our paper, a case-control study in blacks⁴ reported an increased risk with greater than 10 years of use; however, this result was based on an exceedingly small number of controls with long-term exposure ($n = 5$). Estimates based on small sample sizes yield unstable results, because the odds ratio could be markedly affected by the misclassification of only a few subjects.

As suggested by Kundi,¹ we examined the mean (\pm standard deviation) and median years of use in current and

former electric blanket users by menopausal status. Current users had considerably more years of use compared with former users (eg, among postmenopausal controls, mean years of use among current users was 19.5 ± 13.4 compared with 6.5 ± 7.0 in former users). There was no difference in years of use between cases and controls stratified by current/former use. We also looked at the mean age at diagnosis of cases by exposure status and menopausal status. The mean age at diagnosis was not younger for current users or former users compared with never users (eg, among postmenopausal cases, mean age was 64.5 ± 7.2 for current users, 63.1 ± 7.3 for former users, and 62.7 ± 8.1 for never users).

Regarding the laboratory evidence, although there are many experiments designed to study the biologic effects of exposure to ELF-EMF, 2 authoritative reviews have concluded that there is no reproducible evidence of such effects.^{5,6} Even the 3 papers cited by Kundi seem to be at odds as to whether melatonin is involved in the inhibition of cell proliferation. Finally, of the 3 references Kundi cites in support of a specific mechanism, earlier work by 2 of the lead authors has been discredited.^{7,8} Given these considerations, we believe that Kundi's concerns are not well-founded.

Geoffrey C. Kabat*

Erin S. O'Leary

Elinor R. Schoenfeld

M. Cristina Leske

Department of Preventive Medicine

School of Medicine

Stony Brook University

Stony Brook, NY

gckl@optonline.net

Marilie D. Gammon

Department of Epidemiology

School of Public Health

University of North Carolina at Chapel Hill

Chapel Hill, NC

*Affiliation at the time the research was done.

REFERENCES

1. Kundi M. Electric blanket use and breast cancer [Letter]. *Epidemiology* 2004;15:376–377.
2. Kabat GC, O'Leary ES, Schoenfeld ER, et al. Electric blanket use and breast cancer on Long Island. *Epidemiology*. 2003;14:314–320.
3. Laden F, Neas LM, Tolbert PE, et al. Electric blanket use and breast cancer in the Nurses' Health Study. *Am J Epidemiol*. 2000;152:41–49.
4. Zhu K, Hunter S, Payne-Wilkes K, et al. Use of electric bedding devices and risk of breast cancer in African-American women. *Am J Epidemiol*. 2003;158:798–806.
5. National Academy of Sciences. National Research Council. *Possible Health Effects of Exposure to Residential Electric and Magnetic Fields*. National Academy Press; 1997.
6. Portier CJ, Wolfe MS, eds. *National Institute of Environmental Health Sciences Working Group Report Assessment of Health Effects From Exposure to Power-Line Frequency Electric and Magnetic Fields*. NIH Publication No. 98–3981. Research Triangle Park, NC: Research Triangle Park, NC: National Institute of Environmental Health; 1998.
7. Findings of scientific misconduct. *NIH Guide*. June 18, 1999. US Department of Health and Human Services.
8. Adair RK. Measurements described in a paper by Blackman, Blanchard, Benane, and House are statistically invalid. *Bioelectromagnetics*. 1996;17:510–511.

Drinking Water and Cancer

To the Editor:

I very much agree with the commentary¹ of Steenland and Moe, "Epidemiology and Drinking Water, Are We Running Dry?" Their commentary addressed the work of De Roos et al² on the relation between nitrates in public drinking water and the risk of cancer of the colon and rectum. Steenland and Moe suggest that, even given a very favorable milieu within which to conduct a study, these motivated and careful investigators were frustrated in finding results that out-stripped the potential sources of error in their investigation.

Why? Steenland and Moe¹ conclude that the study method—retrospective analysis—is not up to this particular task. They first list the dominant characteristics of the types of problems that are

addressed in studies involving water-borne contaminants. "Exposures are typically low, fairly homogeneous, and hard to measure retrospectively."

Steenland and Moe¹ press on, pointing out that, from a regulatory perspective, this inability to produce results relatively free from uncertainty is worrisome.

The events of September 11, 2001 have produced many repercussions in our society. One that touches public health and the management of water systems is the specter of willful contamination of water supplies. Much has been written about how to systematically respond to such threats.^{4,5} One response has been a surge of innovation in sensor development. Sensors and arrays of sensors are dramatically improving the ability to monitor conditions in the ambient environment for chemical and biologic agents often in real time and at a fraction of the cost of previous technology. An equal challenge for the risk manager is to effectively translate the increased flow of data from such sensor arrays into a meaningful signal with regards to the existence of a possible threat.

Happily, the epidemiologist is not confronted with the same type of pressures as the manager of a large urban water system. The epidemiologist can benefit from the increased precision of integrating exposure to selected contaminants from this new generation of sensors into longitudinal designs.

Conducting prospective studies that benefit from the existence of the new sensor technology would likely reduce the inherent error and uncertainty in measurements taken in the ambient environment or in operational water distribution systems. This seems clear. Why could not the same technologies be adapted for use in exposure assessment at the level of the participant? The analogy which forms in my mind is the benefit which flowed from the miniaturization of air pollution measurement devices and allowed exposure to be quan-

tified in breathing zones and not on the roofs of public buildings.

Does this mean that the application of new technologies will quell all sources of uncertainty of studying water-borne pollutants? No. The investigator must still rely on judgment in carefully designing the study. Some things remain the same.

David M. Gute

Department of Civil and Environmental Engineering
Tufts University
Medford, MA
david.gute@tufts.edu

REFERENCES

1. Steenland K, Moe C. Epidemiology and Drinking Water. Are We Running Dry? *Epidemiology*. 2003;14:635–636.
2. De Roos AJ, Ward MH, Lynch CF, Cantor KP. Nitrate in public water supplies and the risk of colon and rectum cancers. *Epidemiology*. 2003;14:640–649.
3. Ward MH, Cantor KP, Riley D, Merkle S, Lynch CF. Nitrate in public water supplies and risk of bladder cancer. *Epidemiology*. 2003;14:183–190.
4. Beering P. Threats on tap: Understanding the terrorist threat to water. *J Water Resour Plann Manag*. 2002;128:163–167.
5. Haimes Y. Strategic responses to risks of threats to water resources. *J Water Resour Plann Manag*. 2002;128:383–389.

To the Editor:

In the November 2003 issue of EPIDEMIOLOGY, we presented a study¹ on the association between nitrate in public water supplies and the risk of colorectal cancers. An invited commentary on the topic of drinking water studies, written by Drs. Steenland and Moe,² noted the difficulties in assessing drinking water exposures retrospectively, and questioned if epidemiologists could be approaching the limit of what we can learn from retrospective designs. We argue that for studies of drinking water contaminants in relation to cancer, retrospective exposure assessment has often been quite adequate, and alternate study designs have limitations that may preclude them from providing a better solution in the near future.

Despite the challenges of retrospective exposure assessment of drink-

ing water contaminants, these studies have contributed importantly to scientific knowledge about cancer, particularly for arsenic and chlorination byproducts.³ The commentators suggest the prospective study as an improved alternate design because of the possibility of measuring contaminants in individuals' water supplies as well as incorporating intermediate biologic markers which may be relevant for future disease. Studies of arsenic in relation to intermediate biologic effects including micronuclei in urinary bladder cells,⁴ and DNA damage in buccal epithelial cells⁵ have demonstrated the utility of this approach; however, there are few intermediate effects that have been convincingly associated with the development of cancer.⁶ Thus, validation of intermediate markers as cancer precursors must advance considerably before such markers can replace incident cancer as the outcome.

Long-term prospective studies do offer opportunities for more accurate exposure assessment, with the primary advantage being the possibility of sampling the participants' water supplies at the tap, prior to cancer development. This affords a great advantage for participants with private wells, for which few historic monitoring data exist. Nevertheless, the expense of carrying out a long-term prospective study of drinking water contaminants and cancer would necessitate a compelling hypothesis as a rationale for launching the study. Such studies are likely to begin only after substantial evidence accumulates from retrospective epidemiologic studies, supplemented perhaps with a better mechanistic understanding of the carcinogenic process. For these reasons, even when prospective cohort studies have evaluated drinking water contaminants and cancer, they have primarily relied on retrospective exposure data.^{7,8}

Substantial opportunity exists for improving retrospective exposure assessment in drinking water studies.

Models that incorporate improved information on raw water quality and on how treatment processes affect the formation of disinfection byproducts are currently being applied to earlier case-control studies of cancer.⁹ The use of hydraulic simulation models to estimate historic exposure at different points in a water distribution system can also reduce exposure misclassification in retrospective studies.^{10,11} The estimation of historic exposures is likely to be further improved when personal factors affecting exposure^{12,13} and internal dose^{14,15} are used to refine exposure metrics. For example, in our studies of drinking water nitrate and cancer,^{1,16–18} we evaluated subgroups of the population who were likely to have increased formation of N-nitroso compounds.

Drinking water is clearly an essential resource for humans, and epidemiologists should continue to conduct studies to determine what constitutes a "healthful" water supply. Improvement of drinking water studies will depend on further validation of intermediate effect biomarkers as predictors of disease, as well as better exposure assessment through greater knowledge of factors affecting variability in the water supply and in individual exposure and response. Especially as improvements in exposure assessment are applied to retrospective studies, they will continue to play an important role in investigating new hypotheses and testing the consistency of observed associations among different populations.

Anneclaire J. De Roos

Fred Hutchinson Cancer Research Center
Department of Epidemiology
University of Washington
Seattle, WA
aderoos@fhcrc.org

Mary H. Ward

Occupational and Environmental
Epidemiology Branch
Division of Cancer Epidemiology and
Genetics
National Cancer Institute, NIH, DHHS
Rockville, MD

REFERENCES

1. De Roos AJ, Ward MH, Lynch CF, Cantor KP. Nitrate in public water supplies and the risk of colon and rectum cancers. *Epidemiology*. 2003;14:640–649.
2. Steenland K, Moe C. Epidemiology and drinking water: are we running dry? *Epidemiology*. 2003;14:635–636.
3. Cantor KP. Drinking water and cancer. *Cancer Causes Control*. 1997;8:292–308.
4. Biggs ML, Kalman DA, Moore LE, Hopenhayn-Rich C, Smith MT, Smith AH. Relationship of urinary arsenic to intake estimates and a biomarker of effect, bladder cell micronuclei. *Mutat Res*. 1997;386:185–195.
5. Feng Z, Xia Y, Tian D, et al. DNA damage in buccal epithelial cells from individuals chronically exposed to arsenic via drinking water in Inner Mongolia, China. *Anticancer Res*. 2001;21:51–57.
6. Bonassi S, Neri M, Puntoni R. Validation of biomarkers as early predictors of disease. *Mutat Res*. 2001;480–481:349–358.
7. Weyer PJ, Cerhan JR, Kross BC, et al. Municipal drinking water nitrate level and cancer risk in older women: the Iowa Women's Health Study. *Epidemiology*. 2001;12:327–338.
8. Wilkins JR III, Comstock GW. Source of drinking water at home and site-specific cancer incidence in Washington County, Maryland. *Am J Epidemiol*. 1981;114:178–190.
9. American Water Works Association. <http://www.awwarf.com/research/TopicsandProjects/projectSnapshot.aspx?pn=2582>. Accessed November 13, 2003.
10. Gallagher MD, Nuckols J R, Stallones L, Savitz DA. Exposure to trihalomethanes and adverse pregnancy outcomes. *Epidemiology*. 1998;9:484–489.
11. Reif JS, Burch JB, Nuckols JR, Metzger L, Ellington D, Anger WK. Neurobehavioral effects of exposure to trichloroethylene through a municipal water supply. In press: *Environ Res*.
12. Lynberg M, Nuckols JR, Langlois P, et al. Assessing exposure to disinfection by-products in women of reproductive age living in Corpus Christi, Texas, and Cobb county, Georgia: descriptive results and methods. *Environ Health Perspect*. 2001;109:597–604.
13. Miles AM, Singer PC, Ashley DL, et al. Comparison of trihalomethanes in tap water and blood. *Environ Sci Technol*. 2002;36:1692–1698.
14. Mirvish SS, Grandjean AC, Reimers KJ, et al. Effect of ascorbic acid dose taken with a meal on nitrosoproline excretion in subjects ingesting nitrate and proline. *Nutr Cancer*. 1998;31:106–110.
15. Vermeer IT, van Maanen JM. Nitrate exposure and the endogenous formation of carcinogenic nitrosamines in humans. *Rev Environ Health*. 2001;16:105–116.

16. Coss A, Cantor KP, Lynch C, et al. Pancreatic cancer and drinking water and dietary sources of nitrate and nitrite. *Am J Epidemiol*. In press.
17. Ward MH, Mark SD, Cantor KP, Weisenburger DD, Correa-Villasenor A, Zahm SH. Drinking water nitrate and the risk of non-Hodgkin's lymphoma. *Epidemiology*. 1996;7:465–471.
18. Ward MH, Cantor KP, Riley D, Merkle S, Lynch CF. Nitrate in public water supplies and risk of bladder cancer. *Epidemiology*. 2003;14:183–190.

ERRATUM

Kapadia F, Vlahov D, DesJerlais DC, et al. Does bleach disinfection of syringes protect against Hepatitis C infection among young adult injection drug users? *Epidemiology*. 2002;13:738–741.

In Table 1, the percent of cases who never used bleach should be 74%.